

Contourites in the Gulf of Cadiz: a cautionary note on potentially ambiguous indicators of bottom current velocity

Thierry Mulder · Rim Hassan · Emmanuelle Ducassou · Sébastien Zaragosi · Eliane Gonthier · Vincent Hanquiez · Elodie Marchès · Samuel Toucanne

Received: 26 November 2012 / Accepted: 26 May 2013 / Published online: 16 June 2013
© Springer-Verlag Berlin Heidelberg 2013

Abstract Facies associations in cores collected in the deep part of the Gulf of Cadiz, which is under the influence of the lower branch of the Mediterranean Outflow Water, are investigated in terms of the classical contourite model using grain-size analyses and thin sections of indurated sediment. Cores include both low-energy (contourite drift) and high-energy (channel) environments. The thin sections and grain-size distributions show that clayey fine silts and sandy coarse silts are the most common facies associations in the studied contourite sequences, while coarse-grained, gravelly contourites are less common. Grain-size distributions are unimodal in the fine-grained and bi- or trimodal in the coarser-grained contourites. This change in grain-size composition is related both to the partial removal of the fine-grained fraction and to the replenishment of the coarser-grained one. In addition, most of the contacts between individual facies are sharp rather than transitional. This suggests that the contourite sequence is only in part related to changes in bottom current velocity and flow competency, but may also be related to the supply of a coarser terrigenous particle stock, provided by either increased erosion of indurated mud along the flanks of confined contourite channels (mud clasts), or by increased sediment supply by rivers (quartz grains) and downslope mass transport on the continental shelf and upper slope. The classical contourite

facies association may therefore not be solely controlled by current velocity, but may be the product of a variety of depositional histories. The classical contourite depositional sequence should therefore be interpreted with greater care and in the light of the regional sedimentological background. In addition, the wisdom of exclusively using mean or modal particle size for the interpretation of depositional contourite processes is questioned. Instead, it is proposed that the vertical evolution of grain-size populations in the facies successions forming contourite sequences be assessed.

Introduction

The classical contourite sequence depicting a model for deposition from deep-marine contour currents (Hollister 1967) was originally proposed by Gonthier et al. (1984) and Faugères et al. (1984) using data on cores collected from the Faro Drift in the Gulf of Cadiz off the SW Iberian Peninsula (Figs. 1 and 2a). It was first defined for deposits related to sediment transport and deposition by thermohaline circulation or geostrophic currents (Stommel 1958) related to deep-water masses of different temperatures and salinities (Heezen et al. 1966) and known as the “conveyor belt” (Broecker 1991). These deposits form most of the sediment drifts. The contourite concept was extended later to deposits reworked by other deep-sea currents including wind-driven currents, internal tides and other loop currents related to cyclones (see related references in Shanmugan 2012). The model was first applied to clay- and silt-dominated (= muddy) contourites (mud contents of up to 95 %) and later extended to sandy contourite facies by Stow (1994) and used to build a conceptual, idealized contourite sequence model (Stow and Faugères 2008).

The classical scheme comprises two superposed units: a basal coarsening-up unit grading from fine homogeneous

T. Mulder (✉) · R. Hassan · E. Ducassou · S. Zaragosi · E. Gonthier · V. Hanquiez
Université de Bordeaux, UMR CNRS 5805 EPOC,
avenue des facultés,
33405 Talence cedex, France
e-mail: t.mulder@epoc.u-bordeaux1.fr

E. Marchès
Service Hydrographique et Océanographique de la Marine
(SHOM), rue du Chatellier,
29228 Brest cedex 2, France

S. Toucanne
Ifremer, Géosciences Marines, BP 70, 29280 Plouzané, France




Fig. 1 The study area in the Gulf of Cadiz, with an overview of Mediterranean Outflow Water (*MOW*) circulation, the main morphological features (including the Gadiana and Guadalquivir rivers), and the locations of the four cores selected for the study. The bathymetry

(EM 300, Cadisar 1 and 2 cruises) is extracted from Mulder et al. (2003, 2006). *MUW* Mediterranean Upper Water, *MLW* Mediterranean Lower Water, *IMB* Intermediate MOW Branch, *PMB* Principal MOW Branch, *SMB* Southern MOW Branch

mud (clayey fine silt) to mottled coarser silt and finally sandy silt/silty sand, followed by a fining-up unit showing the same facies succession in reverse order (Fig. 2a). This is nicely illustrated in Fig. 2b, which shows a section of a core recovered by T. Mulder and colleagues from the Faro Drift (Cadisar cruise 2, 2004, core CADI2KS08). In the corresponding upcore grain-size distributions (Fig. 2c), curves A and D represent the basal/top mud facies having a typical bimodal shape reflecting the mixing of a subordinate sandy coarse silt and a dominant clayey fine silt population. This bimodal distribution in the terrigenous silt fraction was already underlined by Robinson and McCave (1994), McCave et al. (1995a, b), and Bianchi and McCave (2000). Curves B and C are also bimodal but with a dominant sandy silt/silty sand and a subordinate clayey fine-silt population. This so-called mottled facies corresponds to a sandy mud facies containing coarse silt lenses resulting from infilled polychaete burrows.

Using mean grain size as indicator, the change in sediment composition has hitherto been interpreted as being associated with a change in bottom current competency and velocity. This idea emerged because, differently to what happens in

turbidity currents, contour currents (whatever their origin) act as a winnowing machine generating sediment resuspension by tractive (bed load) transport (Allen 1984) of already deposited particles. The sorting of particles is particularly efficient (Shanmugan 2012). Thus, winnowing intensity and current competency can both be related to current speed, although this concept must be applied with care when dealing with aggregation processes affecting fine-grained cohesive particles. This interpretation is based on the “sortable silt” concept of McCave et al. (1995b), who considered that the non-flocculated fraction can be used as a proxy for current velocity and that the mean (modal) grain size can serve to reconstruct current strength and be interpreted in terms of climate variation (McCave et al. 1995a, b; Bianchi and McCave 1999; Bianchi et al. 2001). This model has been successfully applied to fine-grained sequences examined by Toucanne et al. (2007) in a core collected in the neighbourhood of the present study area, in the low-energy Faro-Cadiz sheeted drift. It shows silt–clay facies (most abundant), clayey silt (mottled) facies and rare clayey sand facies. All the facies have a bimodal grain-size distribution, and are organized into vertical coarsening- and




Fig. 2 Classical contourite sequence. **a** Conceptual scheme (extracted from Gonthier et al. 1984). **b** Core CADI2KS08, Faro Drift (see Fig. 1 for core location): X-ray image and indurated thin sections. **c** Typical grain-size distributions

then fining-up sub-sequences bounded by progressive contacts. They have been interpreted as classical contourites deposited by a current with initially steadily increasing velocity, followed by a decrease in velocity. During periods of increasing current velocity, the coarser mode becomes predominant while the finer diminishes, suggesting significant winnowing of fine particles during periods of enhanced bottom current activity (Hernández-Molina et al. 2003, 2006; Llave et al. 2006).

Stow and Faugères (2008) extended the model to sandy contourites and published a generalized composite contourite model consisting of the superposition of a negatively graded lower sub-sequence (mud + mottled silt and mud units), a middle sandy silt unit and an upper positively graded sub-sequence (mottled silt and mud + mud units). They again underlined the destruction of primary sedimentary structures by thorough bioturbation, and the rarity of sand. As emphasized by Shanmugan (2012), Stow and Faugères (2008) also stated that this idealized conceptual sequence is never found in totality in natural depositional environments and that, commonly, only base- and top-cut sequences with respectively sharper base and top contacts are encountered, reflecting the abrupt change in current velocity or the presence of a complex seafloor topography which locally modifies the current speed. Upper contacts

related to current intensification are more frequent than lower contacts (Hollister 1967).

Although the entire sequence is fully bioturbated, the grain-size signal has evidently been retained, suggesting that it represents longer-term (5,000–20,000 years) fluctuations (acceleration and deceleration) in mean current velocity (Lovell and Stow 1981; Stow et al. 1986) in most terrigenous or mixed drifts. The recurrence intervals of these oscillations might be greater (20,000–40,000 years) in bioclastic accumulations (Stow and Faugères 2008). Such long-term oscillating energy conditions are one of the main differences between contour currents and waning turbidity currents moving downslope along continental margins (Shanmugan 2012). In fact, Faugères and Stow (1993), Stow et al. (2002), and Stow and Faugères (2008) suggest that most of the sandy contourites can be interpreted as bottom current-modified turbidites. Shanmugan (2006) underlined that the predominance of mud in the Faro Drift could be misleading when attempting to define a generalized contourite facies model. Other authors (e.g. Dalrymple and Narbonne 1996) criticised the use of bioturbation as a diagnostic feature of contourites, although most contourites do show intense bioturbation given the low sedimentation rate and the loading with oxygen and nutrients by bottom currents (Chough and Hesse 1985). For example, Tucholke et al.

(1985) prefer to relate the intensity of bioturbation to the current intensity along the seafloor. Masson et al. (2010) showed that the muddy sand contourite collected in the Faroe-Shetland Channel could result from the incomplete winnowing of underlying sand and that along slope transport is minor. Other sedimentary sequences related to contour currents are the gravel-lag contourites resulting from the complete winnowing of particles ranging in size from mud to sand by powerful bottom currents (Stow et al. 1996), resulting in the preservation of the base-only sequence of the generalized Stow and Faugères (2008) model and biogenic contourites (Kidd and Hill 1986).

Other studies close to the Gulf of Cadiz—for example, on the Galicia margin (e.g. Bender et al. 2012) or in the Mediterranean (e.g. Verdicchio and Trincardi 2008)—have provided evidence for recent (post-Last Glacial Maximum) sedimentation on continental slopes resulting from increased sediment availability (including variations in detrital input, marine productivity and sea-level stands, as well as current velocities). Contourite deposition, however, has been more strongly affected by climatic and sea-level changes rather than tectonics since the Mid-Pleistocene Revolution (Llave et al. 2011). The formation of coarse-grained contourites in the Gulf of Cadiz over the last 50,000 years is thus interpreted to be the result of an intensification of MOW formation and current velocity during cold periods. Sandy-silt contourites are related to the Younger Dryas, Heinrich events (in particular H1 and H2) and stadial Dansgaard-Oeschger events (Mulder et al. 2002; Hanquiez 2006; Llave et al. 2006; Voelker et al. 2006; Toucanne et al. 2007). Conversely, fine-grained contourites are related to warm periods (Bølling-Allerød and Dansgaard-Oeschger interstadials). Such fluctuations in the MOW and other bottom currents, and their implication for late Quaternary climate forcing on slope sedimentation, have also been recorded in the northernmost areas of MOW activity, i.e. along the Rockall Trough and Porcupine Bank, and off Ireland (Øvrebø et al. 2005, 2006). However, an increase in bottom current velocity during sea-level lowstands is not a general rule, but also depends on regional tectonic activity and the presence of a nearby englacial continental margin (Brackenridge et al. 2011).

The fact that the Gulf of Cadiz is located close to the outflow of the Strait of Gibraltar (Fig. 1) makes it a prime target for the study of contourites, i.e. sedimentary beds deposited by bottom currents following bathymetric contours. Within this context, the main aim of the present paper is to call for caution in interpreting the classical contourite model characterized by a coarsening- and fining-up depositional sequence solely in terms of changes in current velocity because other factors can also explain changes in the grain-size composition of the sediment. It is also demonstrated that mean grain size alone is insufficient for the reconstruction of

contourite formation mechanisms and that the temporal (vertical) evolution of the whole grain-size distributions yields much more information for comprehensive interpretations.

Physical setting

The continental margin of the Gulf of Cadiz is exposed to the action of a strong, warm (13 °C) and saline ($>37 \text{ g } \Gamma^{-1}$) current called the Mediterranean Outflow Water (MOW), which forms in the Alboran Sea (western Mediterranean) and flows into the North Atlantic Ocean (Fig. 1). In detail, the MOW is formed by the superposition of two major flow systems: the Mediterranean Upper Water (MUW; salinity $36.1 \text{ g } \Gamma^{-1}$, temperature 11.5 °C; Ambar 1983), which forms the uppermost core funnelled through the Alvarez Cabral Channel, and the Mediterranean Lower Water (MLW; salinity $37.4 \text{ g } \Gamma^{-1}$, temperature 13.6 °C; Ambar 1983), which comprises three branches—the Intermediate Branch (IMB), which flows between 700 and 900 m water depth, the Principal Branch (PMB) between 900 and 1,000 m water depth, and the Southern Branch (SMB) between 1,000 and 1,200 m water depth. Just after the Strait of Gibraltar is passed, part of the MOW is sharply deflected towards the Spanish and Portuguese margins under the effect of the Coriolis force (Madelain 1970). When flowing through the Strait of Gibraltar the velocity of the MOW exceeds 2 m s^{-1} (Fig. 1). Although the mean velocity then rapidly decreases, obstacles can locally accelerate the current along the seabed (García et al. 2009). On average, the velocity reaches only 0.2 m s^{-1} off Cape St. Vincent (southwest Portugal), where the MOW lifts off the seafloor at about 1,400 m water depth.

The bathymetric map (Fig. 1) shows that the seafloor morphology in the eastern part of the Gulf of Cadiz is strongly affected by the MOW. In addition, the regional sedimentation patterns have been controlled by the MOW since Messinian times, and the long depositional history therefore excellently reveals the general sedimentary regime in the Gulf of Cadiz. The resulting sedimentary system has been called the Cadiz contourite depositional system (CDS) by Hernández-Molina et al. (2003, 2011). The MOW captures particles supplied by Spanish rivers and also erodes material from the continental shelf (Grousset et al. 1988). Seafloor morphology strongly influences the circulation pattern of the MOW, as is also known from other contourite depositional systems such as those in the Mediterranean (e.g. Palomino et al. 2011). Close to Gibraltar, a gravel lag and large-scale erosional structures (furrows) have been observed (Kenyon and Belderson 1973). Downflow, the deposits become silty with sand patches and sand waves, which is in agreement with the progressive decrease in MOW energy (Kenyon and Belderson 1973; Mélières 1974;

Faugères et al. 1985a; Nelson et al. 1993, 1999). The sedimentary bodies, which occur mainly in valleys between adjacent ridges through which the MOW is channelized, show a progressive downflow decrease in grain size. On the seaward side of the channels, the MOW velocity decreases sharply and fine particles are deposited to form thick successive sediment accumulations, so-called contourite drifts (Mougenot and Vanney 1982). Of these, the Faro, Albufeira, Portimao and Lagos drifts are formed by the MUW, whereas the Guadalquivir Drift is formed by the MLW (Llave et al. 2007; García et al. 2009). The drifts are composed of muddy sediments and grew in size from the Messinian to the Present after the Mediterranean–Atlantic connection was formed (Faugères et al. 1985b). Despite the presence of erosional surfaces within the drift sequences (Llave et al. 2006), deposition in the contourite drifts is mostly continuous, thus enabling good palaeoceanographic reconstructions (e.g. Voelker et al. 2006; Toucanne et al. 2007). It is for this reason that the original contourite type sequence was described from this region (Gonthier et al. 1984; core CADI2KS08, Figs. 1 and 2). In the western part of the Gulf of Cadiz, the energy of the MOW is low and sedimentation therefore dominated by downslope gravity processes responsible for creating the classical seafloor morphology characterized by canyons and gullies.

Another part of the MOW flows westwards and, after spilling over a topographic high, strongly decelerates to deposit its load in the form of extensive fields of sediment waves in the 1,000–1,200 m depth interval (Fig. 1). This flow may be partly channelled through large conduits such as the Gil Eanes Channel. The deceleration after spillover results in high sedimentation rates forming the giant contouritic levee identified by Mulder et al. (2003). These high sedimentation rates in association with active seismicity in the Gulf of Cadiz and neighbouring areas (Zitellini et al. 1999; Gutscher et al. 2002) have generated numerous slope failures on the leeward side of the levee (Mulder et al. 2009).

Material and methods

This study is based on grain-size analyses and indurated thin sections of four Kullenberg cores collected along the pathway of the MLW during two cruises aboard the RV *Suroit* (see Fig. 1 for locations): core CADI2KS08 (cf. above, Cadisar 2 cruise, September 2004), and cores CADKS14, 18 and 24 (Cadisar 1 cruise, August 2001). Mulder et al. (2003, 2006) have provided some preliminary information on the physical setting characterizing these cores.

The typical contourite sequence (Fig. 2) sampled in core CADI2KS08 (36°40.890'N, 08°06.240'W, water depth 789 m) is from the sheeted part of the Faro Drift affected by the MLW/IMB. This sequence is dated from the Younger Dryas

(Gonthier et al. 2010). Core CADKS14 (36°24.998'N, 07°31.044'W, water depth 738 m) is located on the Guadalquivir Drift close to the Guadalquivir Channel affected by the MLW/PMB. The studied sequence in this core is dated from Marine Isotopic Stage 3 (Gonthier et al. 2010). It should represent classical sedimentation within a contourite drift, similar to core CADI2KS08 from the Faro Drift. It was chosen for this study because it is affected by the same water mass as cores CADKS18 and CADKS24, i.e. the MLW.

Core CADKS18 (36°18.08'N, 07°24.78'W, water depth 1,001 m) is located southeast of the Guadalquivir diapiric ridge, in a circular topographic low interpreted as a dissolution depression, and is affected by the MLW/SMB. No clear dating could be obtained on this core because of intense reworking. Core CADKS24 (36°04.944'N, 07°56.523'W, water depth 1,316 m) is located in the very distal part of the Cadiz CDS, in the south-western sector of the Cadiz Channel, and also affected by the MLW/SMB. The thin section has been deposited during the Last Glacial Maximum (Gonthier et al. 2010). Core interpretation shows the superposition of muddy contourites with silty sand intercalations.

Indurated 30- μ m thin sections of soft sediment were obtained according to the impregnation protocol described by Zaragosi et al. (2006). Fresh core samples were dehydrated sequentially in acetone, and hardened with an epoxy resin. Fluorescent dye was added to the resin to enable subsequent fluorescent light analysis by which indurated and non-indurated parts can be distinguished. The bonded blocks were cut to approximately 100 μ m using a precision saw and thereafter hand polished to a thickness of 30 μ m by means of a rotating lapidary unit. Thin-section images were acquired under a fully automated Leica DM6000 B digital microscope at multiple magnifications.

Bulk sediment subsamples were taken for grain-size measurements using a Malvern MasterSizer S (0.05 to 878.67 μ m). Mud is defined as the <63 μ m fraction, i.e. silt and clay (Twenhofel 1937; cf. also Flemming 2000).

Results

In the selected cores, three contourite facies are observed: muddy contourites (Fig. 3, core CADKS24), silty contourites (Figs. 4 and 5, cores CADKS14 and CADKS18) and gravelly contourites (Fig. 5, core CADKS18). Muddy and silty–sandy contourites represent the typical facies of the base/top and middle contourite sequences respectively (cf. Fig. 2, core CADI2KS08). Gravelly contourites represent a more unusual facies in higher-energy environments (Stow and Faugères 2008). An intermediate facies composed of sand is not represented in the available data. Such a facies would represent lag deposits in contourite channels (Hernández-Molina et al. 2003; Hanquiez et al. 2007). Sandy contourites also form




Fig. 3 Core CADKS24, distal sector of Cadiz contourite depositional system. **a** Core log (*c* clay, *s* silt, *fs* fine sand, *ms* medium sand, *cs* coarse sand; see Fig. 1 for core location). **b** Muddy contourite: X-ray, grain size, and indurated thin sections under natural light and fluorescence

cm-thick beds in cores recovered from essentially fine-grained contourite drifts (Gonthier et al. 1984) and correspond to periods of increased MOW velocity (colder climate periods) or represent coarse-grained deposits within contourite channels mostly in the so-called ridge and channel sector of the CDS (Hernández-Molina et al. 2003; Hanquiez et al. 2007, 2010). These would be interpreted in a similar way as the gravelly contourite facies illustrated in this paper.

Most of the deposits in contourite drifts are composed of muddy sediment (Fig. 3; Gonthier et al. 1984). This is also the most common facies found in the classical contourite sequence (Fig. 2). The muddy contourites are composed of clay and fine silt with subordinate contributions of coarse silt and sand. Grain-size analyses (Fig. 3) show distributions with a high peak located in the clay–fine silt range extending from 3 to 30 μm .

The silty contourites (Fig. 4) show a vertical superposition of clayey fine silts alternating with coarse silt–fine sand, the former corresponding to the so-called mottled facies. Thin sections clearly show that the transitions between the two

facies are sharp rather than continuous (*c1* and *c2* in Fig. 4). Grain-size measurements along the thin sections confirm the sharp transition between the mottled mud and the coarse silt–fine sand facies. Below the sharp *c1* contact, the quartz particles have a modal diameter of about 10 μm (fine silt), whereas above the sharp *c1* contact the grain-size distribution curves are bimodal with a second peak around 40 μm (medium to coarse silt), thus revealing the arrival of a coarser quartz silt population at this study site. This change in sediment composition corresponds to the transition from the fine-grained mud facies to the muds containing silt lenses within the mottled facies. The silt deposits are laminated, the laminations being enhanced by several millimetre-long elongated clasts of mollusc shells (*ms* in Fig. 4). Above the sharp *c2* contact, the grain-size distribution becomes trimodal, signifying the arrival of a third particle group having modal diameters ranging from 100 to 110 μm . This population also consists of quartz grains and includes a few indurated sand-sized mud clasts.




Fig. 4 Core CADKS14, Guadalquivir Drift. **a** Core log (see Fig. 1 for core location and Fig. 3 for legend). **b** Silty contourite: X-ray, grain size, and indurated thin sections under natural light and fluorescence; *c1*, *c2* sharp contacts, *s* sand, *ms* mollusc shell, *cmc* consolidated mud clasts




Fig. 5 Core CADKS18, topographic low near Guadalquivir Ridge. **a** Core log (see Fig. 1 for core location and Fig. 3 for legend). **b** Laminated and gravely contourite (lower right panel extracted from Faugères and Mulder 2011): X-ray, grain size, and indurated thin sections under natural light and fluorescence; *c* contact, *lf* laminated facies, *cmc* consolidated mud clasts

Figure 5 shows a thin section with a similar superposition of several alternating laminae composed of mud and very fine sand. In these silt-dominated contourites, the grain-size distribution is bimodal, the second particle group having a well-defined modal diameter $>110\ \mu\text{m}$ (very fine–fine sand). It is composed of quartz and a few sand-sized indurated mud clasts.

In addition to the change in grain-size composition, the fluorescence analysis revealed a change in the nature of the grains below and above the sharp contacts (Figs. 3, 4 and 5). In the muddy facies below, the dark grains in the fluorescence image consist entirely of quartz. In the silty facies above, by contrast, the dark grains are formed by quartz and sand/silt-sized indurated mud clasts, suggesting that consolidated mud has been eroded along the path of the MOW and subsequently redeposited in the contourites. This is an indication that the facies change corresponds to an increase in flow velocity/competency accompanied by a new input of one or more grain-size populations derived from either an autochthonous source represented by bottom sediment erosion (indurated sand/silt-sized mud clasts) or an allochthonous (external or remote) source supplying sand-sized quartz particles (e.g. by slope failure and turbidite activity, increased river discharge). For example, Rosa et al. (2011) have shown that, during periods of sea-level lowstands, the Guadiana River discharges its bed load directly onto the SW Iberia Shelf. During sea-level highstands, by contrast, the bed load drops out at the river mouth, whereas the suspended load is carried out to sea.

Figure 5 also shows a thin section with a sharp contact between a silty contourite (coarse silt and fine sand) and a gravelly contourite. In the very coarse part above the sharp contact, mm-size grains float in a sandy matrix without showing any preferred grain orientation. Three grain-size populations are observed. The grain-size distributions of the two finest ones have been measured by means of a Malvern microgranulometer and are visible as discrete peaks on the grain-size curves. The coarsest grain-size population is made of aggregated and slightly lithified (darker on the fluorescence image) mud clasts which are visible only in the thin section. This population is still associated with fine grains but in much lower concentrations compared to the mud facies.

Discussion

Thin section analysis shows that most of the contacts between the classical contourite facies (mottled, fine sand, coarse sand) are sharp. This suggests that the facies transitions do not reflect a steady increase in current competency but rather the response to sudden events, consistent with suggestions by Stow and Faugères (2008). The contacts may indicate periods of sediment bypassing with eventual erosion of the bed itself.

In most cases the clayey fine silt population ($5\text{--}8\ \mu\text{m}$) remains present in all the facies (Fig. 3). Consequently, the increase of the mean grain size in contourite facies is not necessarily the result of winnowing of the finest fraction but possibly a consequence of the supply of additional coarser silt particles which change the proportional composition in favour of the coarser fraction. The presence of laminations enhanced by shell fragments and the concentration of quartz grains suggest a discrete grain (sortable silt) input as bed load at the base of the contour current (cf. McCave 2008; Masson et al. 2010). This confirms that bed load transport is a major characteristic of contour current deposition (Shanmugan 2012).

In other cases (Fig. 4b), the grain-size evolution in the superposed facies forming the contourite sequence suggests either a progressive removal of the clayey fine silt population, and a corresponding proportional enhancement of the coarser silt population without the input of new material, or simply a replenishment of the coarser silt population with additional material resulting in its proportional growth relative to the finer population. But once again, the sharp transition between facies suggests that this replacement is not a continuous process.

This interpretation—mud and sand are both transported and deposited by contour currents during periods of increased current velocity combined with increased sand particle supply—is consistent with the hypothesis of Masson et al. (2010). This is also consistent with the clean sand concentrations observed in all channels of the Gulf of Cadiz (Hanquiez et al. 2007), and explains the better sorting of sandy contourites when compared to sandy turbidites as suggested by Shanmugan (2012).

Classical interpretations using Hjulström- or Shields-type diagrams suggest that a progressive increase in flow velocity is necessary to erode and transport particles corresponding to non-cohesive clay and fine silt, coarse silt, fine sand, cohesive mud and coarse sand, in that order. This is consistent with the classical interpretation of the increase in the mean grain size in contourite sequences. However, the fact that the increase in mean grain size can result from both the removal of the finer fraction and/or a new supply of coarser grains suggests that contourite sequences cannot only be interpreted as simply reflecting an increase in flow competency. A slight increase in flow velocity can certainly be responsible for the progressive removal of the finest grains if these are non-cohesive. In addition, a more substantial increase in current velocity may involve the mobilisation and downstream transport of “new” particles which may consist (1) of autochthonous indurated mud clasts eroded from upstream contourite channel flanks, and/or (2) of allochthonous sand- and silt-sized quartz grains derived from more remote (external) sources.

In areas where the current velocity is low, eroded mud clasts would be mixed with the pre-existing particles. The laminated facies (If in Fig. 5, core CADKS18) suggests that

the coarse particle stock is mobilized or supplied rhythmically within the period of contourite deposition. This rhythm is probably masked by the bioturbation in most of the contourite sequences, as represented in particular by the mottled facies where primary structures are rarely preserved. In the case of core CADKS18, however, primary sedimentary structures are preserved because of a lack of bioturbation and erosion, consistent with the core being located in a topographic depression. Although this situation is quite rare, it nevertheless contributes towards a better understanding of how contourite sequences can be generated.

Conclusions

The classical model of a contourite sequence described from the deeper parts of the Gulf of Cadiz, and which has been interpreted as the response to a successive increase and decrease in contour current velocity and competency, is here shown to be potentially more complex than that. This paper also shows that the vertical grain-size evolution and the intensity of bioturbation are good diagnostic criteria for the recognition of contourites. Detailed analyses of grain-size distributions and the identification of grain-size populations have thus revealed a number of additional potential sedimentary processes involved in contourite deposition.

1. An increase in flow velocity can result in the progressive removal of the finest particles, while rhythmic velocity pulses produce alternations of silt- and clay-dominated mud laminae.
2. Rare preserved laminae confirm the importance of bed load transport in contourite deposition.
3. The finer/coarser grain alternations and all primary sedimentary structures can be destroyed by bioturbation, making process–response interpretations difficult but indicate that the deposition of the whole sequence has taken a considerable length of time.
4. Increases in current velocity result in periods of sediment bypassing which are recorded by sharp contacts at the top of the underlying mud facies; these sharp contacts are enhanced by a new supply of coarse particles which are derived either from the erosion of local upstream deposits or from an input of sediment from external sources (e.g. downslope processes).
5. With increasing MOW velocity and, hence, competency, the same facies succession can be derived by either the selective removal of the finer fraction or a supply of new sediment which may be eroded from an upstream autochthonous source—e.g. the side walls of contourite channels (indurated muddy clasts)—and/or an external allochthonous source supplying a new stock of coarser-grained quartz derived from remote sources such as high river loads during cold periods. Either process enhances the proportional contribution of one population relative to the other.
6. The concept of sortable silt can be used in interpreting grain-size distributions in contourites but should be associated with a determination of the nature of the grains and/or clasts constituting the deposits.

Acknowledgements We thank the crew of the RV *Le Suroît* for technical assistance during the Cadisar 1 and 2 cruises, and F.J. Hernández-Molina and R.G. Lucchi for reviews of the manuscript. We are grateful to B. Martin for the realisation of thin sections on indurated sediments. We thank the editors of Geo-Marine Letters for editorial and language correction on the manuscript.

References

- Allen JRL (1984) Sedimentary structures: their character and physical basis, vol I. Elsevier, Amsterdam
- Ambar I (1983) A shallow core of Mediterranean water off western Portugal. *Deep Sea Res A* 30(6):677–680
- Bender VB, Hanebuth TJJ, Mena A, Baumann K-H, Francés G, von Döbenek T (2012) Control of sediment supply, palaeoceanography and morphology on late Quaternary sediment dynamics at the Galician continental slope. *Geo-Mar Lett* 32:313–335
- Bianchi GG, McCave IN (1999) Holocene periodicity in North Atlantic climate and deep-ocean flow south of Iceland. *Nature* 397:515–517
- Bianchi GG, McCave IN (2000) Hydrography and sedimentation under the deep western boundary current on Björn and Gardar Drifts, Iceland Basin. *Mar Geol* 165:137–169
- Bianchi GG, Vautravers M, Shackleton NJ (2001) Deep flow variability under apparently stable North Atlantic Deep Water production during the last interglacial of the subtropical NW Atlantic. *Paleoceanography* 16:306–316
- Brackenridge R, Stow DAV, Hernández-Molina FJ (2011) Contourites within a deep-water sequence stratigraphic framework. *Geo-Mar Lett* 31:343–360
- Broecker WS (1991) The great ocean conveyor. *Oceanography* 4:79–89
- Chough SK, Hesse R (1985) Contourites from the Eirik Ridge, South of Greenland. *Sediment Geol* 41:185–199
- Dalrymple RW, Narbonne GM (1996) Continental slope sedimentation in the Sheepbed Formation (Neoproterozoic, Windermere Supergroup), Mackenzie Mountains, N.W.T. *Can J Earth Sci* 33:848–862
- Faugères J-C, Mulder T (2011) Contour currents and contourite drifts. In: Hüeneke H, Mulder T (eds) *Deep-sea sediments*, vol 63, *Developments in Sedimentology*. Elsevier, Amsterdam, pp 149–214
- Faugères J-C, Stow DAV (1993) Bottom-current-controlled sedimentation: a synthesis of the contourite problem. *Sediment Geol* 82:287–297
- Faugères J-C, Gonthier E, Stow DAV (1984) Contourite drift molded by deep Mediterranean outflow. *Geology* 12:296–300
- Faugères J-C, Frappa M, Gonthier E, Grousset FE (1985a) Impact de la veine d'eau méditerranéenne sur la sédimentation de la marge sud et ouest ibérique au Quaternaire récent. *Bull Inst Géol Bassin Aquitaine* 37:259–287
- Faugères J-C, Cremer M, Monteiro H, Gaspar L (1985b) Essai de reconstitution des processus d'édification de la ride sédimentaire

- du Faro (marge sud-portugaise). *Bull Inst Géol Bassin Aquitaine* 37:229–258
- Flemming BW (2000) A revised textural classification of gravel-free muddy sediment on the basis of ternary diagrams. *Cont Shelf Res* 20(10):1125–1137
- García M, Hernández-Molina FJ, Llave E, Stow DAV, León R, Fernández-Puga MC, Díaz del Río V, Somoza L (2009) Contourite erosive features caused by the Mediterranean Outflow Water in the Gulf of Cadiz: Quaternary tectonic and oceanographic implications. *Mar Geol* 257:24–40
- Gonthier E, Faugères J-C, Stow DAV (1984) Contourite facies of the Faro Drift, Gulf of Cadiz. In: Stow DAV, Piper DJW (eds) *Fine-grained sediments: deep-water processes and facies*. Geological Society by Blackwell Scientific Publications Oxford London Edinburgh Boston Palo Alto Melbourne Spec Publ 15:275–292
- Gonthier E, Marchès E, Hanquiez V, Hassan R, Ducassou E, Mulder T, Zaragosi S (2010) Influence of sea-level and climate on the contourite deposits in the Gulf of Cádiz. In: *Proc Congr Deep-water Circulation: Processes & Products*, vol 7. Geo-Temas, Baiona, pp 53–54
- Grousset FE, Joron JL, Biscaye PE, Latouche C, Treuil M, Maillet N, Faugères JC, Gonthier E (1988) Mediterranean outflow through the Strait of Gibraltar since 18,000 years B.P.: mineralogical and geochemical arguments. *Geo-Mar Lett* 8:25–34
- Gutscher M-A, Malod J, Rehault J-P, Contrucci I, Klingelhoefer F, Mendes-Victor L, Spakman W (2002) Evidence for active subduction beneath Gibraltar. *Geology* 30(12):1071–1074
- Hanquiez V (2006) *Processus sédimentaires et activité de la Veine d'Eau Méditerranéenne au cours du Quaternaire terminal dans le Golfe de Cadix*. PhD Thesis, N° 3333, Université Bordeaux, p 352
- Hanquiez V, Mulder T, Lecroart P, Gonthier E, Marchès E, Voisset M (2007) High resolution seafloor images in the Gulf of Cadiz, Iberian margin. *Mar Geol* 246:42–59
- Hanquiez V, Mulder T, Toucanne S, Lecroart P, Bonnel C, Marchès E, Gonthier E (2010) The sandy channel-lobe depositional systems in the Gulf of Cadiz: gravity processes forced by contour current processes. *Sediment Geol* 229:110–123
- Heezen BC, Hollister CD, Ruddiman WF (1966) Shaping of the continental rise by deep geostrophic contour currents. *Science* 152:502–508
- Hernández-Molina FJ, Llave E, Somoza L, Fernández-Puga MC, Maestro A, León R, Barnolas A, Medialdea T, García M, Vázquez JT, Díaz del Río V, Fernández-Salas LM, Lobo F, Alveirinho Dias JM, Rodero JY, Gardner J (2003) Looking for clues to paleoceanographic imprints: a diagnosis of the Gulf of Cadiz contourite depositional systems. *Geology* 31(1):19–22
- Hernández-Molina FJ, Llave E, Stow DAV, García M, Somoza L, Vázquez JT, Lobo FJ, Maestro A, Díaz del Río V, León R (2006) The contourite depositional system of the Gulf of Cadiz: a sedimentary model related to the bottom current activity of the Mediterranean outflow water and its interaction with the continental margin. *Deep-Sea Res II Top Stud Oceanogr* 53(11/13):1420–1463
- Hernández-Molina FJ, Serra N, Stow DAV, Llave E, Ercilla G, Van Rooij D (2011) Along-slope oceanographic processes and sedimentary products around the Iberian margin. *Geo-Mar Lett* 31:315–341
- Hollister CD (1967) *Sediment distribution and deep circulation in the western North Atlantic*. PhD dissertation, Columbia University, New York
- Kenyon NH, Belderson RH (1973) Bed-forms of the Mediterranean undercurrent observed with side-scan sonar. *Sediment Geol* 9:77–99
- Kidd RB, Hill P (1986) Sedimentation on mid-ocean sediment drifts. In: Summerhayes CP, Shackleton NJ (eds) *North Atlantic Paleooceanography*. *Geol Soc London Spec Publ.*, pp 87–102
- Llave E, Schönfeld J, Hernández-Molina FJ, Mulder T, Somoza L, Díaz del Río V, Sánchez-Almazo I (2006) High-resolution stratigraphy of the Mediterranean outflow contourite system in the Gulf of Cadiz during the late Pleistocene: the impact of Heinrich events. *Mar Geol* 227(3/4):241–262
- Llave E, Hernández-Molina FJ, Somoza L, Stow DAV, Díaz Del Río V (2007) Quaternary evolution of the contourite depositional system in the Gulf of Cádiz. *Geol Soc Lond Spec Publ* 276:49–79
- Llave E, Matias H, Hernández-Molina FJ, Ercilla G, Stow DAV, Medialdea T (2011) Pliocene–Quaternary contourites along the northern Gulf of Cadiz margin: sedimentary stacking pattern and regional distribution. *Geo-Mar Lett* 31:377–390
- Lovell JPB, Stow DAV (1981) Identification of ancient sandy contourites. *Geology* 9:347–349
- Madelain F (1970) Influence de la topographie du fond sur l'écoulement méditerranéen entre le détroit de Gibraltar et le Cap Saint Vincent. *Cah Oceanogr* 22:43–61
- Masson DG, Plets RMK, Huvenne VAI, Wynn RB, Bett BJ (2010) Sedimentology and depositional history of Holocene sandy contourites on the lower slope of the Faroe-Shetland Channel, northwest of the UK. *Mar Geol* 268:85–96
- McCave IN (2008) Size sorting during transport and deposition of fine sediments: sortable silt and flow speed. In: Rebesco M, Camerlenghi A (eds) *Contourites*, vol 60, *Developments in Sedimentology*. Elsevier, Amsterdam, pp 121–142
- McCave IN, Manighetti B, Beveridge NAS (1995a) Circulation in the glacial North Atlantic inferred from grain-size measurements. *Nature* 374:149–152
- McCave IN, Manighetti B, Robinson SG (1995b) Sortable silt and fine sediment size/composition slicing: parameters for palaeocurrent speed and palaeoceanography. *Paleoceanography* 10:593–610
- Mélières F (1974) *Recherches sur la dynamique sédimentaire du Golfe de Cadix (Espagne)*. Thèse Sci. Nat., Paris VI, p 1–235
- Mougenot D, Vanney JR (1982) Les rides de contourites plio-quaternaires de la pente continentale sud-portugaise. *Bull Inst Géol Bassin Aquitaine* 3:131–139
- Mulder T, Lecroart P, Voisset M, Schönfeld J, Le Drezen E, Gonthier E, Hanquiez V, Zahn R, Faugères J-C, Hernández-Molina FJ, Cadisar scientific party (2002) Past deep-ocean circulation and the paleoclimate record - Gulf of Cadiz. *EOS Am Geophys Union Trans* 83(43):481–488
- Mulder T, Voisset M, Lecroart P, Le Drezen E, Gonthier E, Hanquiez V, Faugères J-C, Habgood E, Hernández-Molina FJ, Estrada F, Llave-Barranco E, Poirier D, Gorini C, Fuchey Y, Voelker A, Freitas P, Lobo Sanchez F, Fernandez LM, Kenyon NH, Morel J (2003) The Gulf of Cadiz: an unstable giant contouritic levee. *Geo-Mar Lett* 23:7–18
- Mulder T, Lecroart P, Hanquiez V, Marches E, Gonthier E, Guedes J-C, Thiébot E, Jaaidi B, Kenyon N, Voisset M, Perez C, Sayago M, Fuchey Y, Bujan S (2006) The western part of the Gulf of Cadiz: contour currents and turbidity currents interactions. *Geo-Mar Lett* 26:31–41
- Mulder T, Gonthier E, Lecroart P, Hanquiez V, Marchès E, Voisset M (2009) Sediment failures and flows in the Gulf of Cadiz (eastern Atlantic). *Mar Petrol Geol* 26:660–672
- Nelson CH, Baraza J, Maldonado A (1993) Mediterranean undercurrent sandy contourites, Gulf of Cadiz, Spain. In: Stow DAV, Faugères J-C (eds) *Contourites and bottom currents. Contourites and Hemipelagites in the Deep-Sea*. *Sed Geol* 82:103–131
- Nelson CH, Baraza J, Maldonado A, Rodero J, Escutia C, Barber JH Jr (1999) Influence of the Atlantic inflow and Mediterranean outflow currents on Quaternary sedimentary facies of the Gulf of Cadiz continental margin. In: Maldonado A, Nelson CH (eds) *Evolution of the Iberian Margin and the Gulf of Cadiz*. *Mar Geol* 155:99–129
- Øvrebo LK, Haughton PDW, Shannon PM (2005) Temporal and spatial variations in Late Quaternary slope sedimentation along the undersupplied margins of the Rockall Trough, offshore west Ireland. *Nor J Geol* 85:279–294

- Øvrebø LK, Haughton PDW, Shannon PM (2006) A record of fluctuating bottom currents on the slopes west of the Porcupine Bank, offshore Ireland – implications for late Quaternary climate forcing. *Mar Geol* 255:279–309
- Palomino D, Vázquez J-T, Ercilla G, Alonso B, López-González N, Díaz-del-Río V (2011) Interaction between seabed morphology and water masses around the seamounts on the Motril Marginal Plateau (Alboran Sea, Western Mediterranean). *Geo-Mar Lett* 31:465–479
- Robinson SG, McCave IN (1994) Orbital forcing of bottom-current enhanced sedimentation on Feni Drift, NE Atlantic, during the mid-Pleistocene. *Paleoceanography* 9:943–972
- Rosa F, Dias JA, Mendes I, Ferreira Ó (2011) Mid to late Holocene constraints for continental shelf mud deposition in association with river input: the Guadiana Mud Patch (SW Iberia). *Geo-Mar Lett* 31:109–121
- Shanmugan G (2006) Deep-water processes and facies models: implications for sandstone petroleum reservoirs, vol 5, *Handbook of Petroleum Exploration and Production*. Elsevier, Amsterdam
- Shanmugan G (2012) New perspectives on deep-water sandstones: origin, recognition, initiation and reservoir quality, vol 9, *Handbook of Petroleum Exploration and Production*. Elsevier, Amsterdam
- Stommel H (1958) The abyssal circulation. *Deep-Sea Res* 5:80–82
- Stow DAV (1994) Deep-sea processes of sediment transport and deposition. In: Pye K (ed) *Sediment transport and depositional processes*. Blackwell, Oxford, pp 257–291
- Stow DAV, Faugères J-C (2008) Contourite facies and the facies model. In: Rebesco M, Camerlenghi A (eds) *Contourites*, vol 60, *Developments in Sedimentology*. Elsevier, Amsterdam, pp 223–256
- Stow DAV, Faugères J-C, Gonthier E (1986) Facies distribution and drift growth during the late Quaternary (Gulf of Cadiz). *Mar Geol* 72:71–100
- Stow DAV, Reading HG, Collinson J (1996) Deep seas. In: Reading HG (ed) *Sedimentary environments*, 3rd edn. Blackwell, Oxford, pp 395–453
- Stow DAV, Faugères J-C, Howe JA, Pudsey CJ, Viana AR (2002) Bottom currents, contourites and deep-sea sediment drifts: current state-of-the-art. In: Stow DAV, Pudsey CJ, Howe JA, Faugères J-C, Viana AR (eds) *Deep-water contourite systems: modern drifts and ancient series, seismic and sedimentary characteristics*. *Geol Soc London Spec Publ* 22:7–20
- Toucanne S, Mulder T, Schönfeld J, Hanquiez V, Gonthier E, Duprat J, Cremer M, Zaragosi S (2007) Contourites of the Gulf of Cadiz: a high-resolution record of the paleocirculation of the Mediterranean outflow water during the last 50,000 years. *Palaeogeogr Palaeoclimatol Palaeoecol* 246:354–366
- Tucholke BE, Hollister CD, Biscaye PE, Gardner WD (1985) Abyssal current character determined from sediment bedforms on Nova Scotian continental rise. *Mar Geol* 66:43–57
- Twenhofel WH (1937) Terminology of the fine-grained mechanical sediments. National Research Council, Report of the Commission on Sedimentation for 1936–1937, pp 81–104
- Verdicchio G, Trincardi F (2008) Mediterranean shelf-edge muddy contourites: examples from the Gela and South Adriatic basins. *Geo-Mar Lett* 28:137–151
- Voelker AHL, Lebreiro SM, Schönfeld J, Cacho I, Erlenkeuser H, Abrantes F (2006) Mediterranean outflow strengthening during northern hemisphere coolings: a salt source for the glacial Atlantic? *Earth Planet Sci Lett* 245:39–55
- Zaragosi S, Bourillet J-F, Eynaud F, Toucanne S, Denhard B, Van Toer A, Lanfumey V (2006) The impact of the last European deglaciation on the deep-sea turbidite systems of the Celtic-Armorican margin (Bay of Biscay). *Geo-Mar Lett* 26(6):317–329
- Zitellini N, Chierici F, Sartori R, Torelli L (1999) The tectonic source of the 1755 Lisbon earthquake and tsunamis. *Ann Geofis* 42:49–55